

Electroproduction of vector mesons off nuclei at ELFE : exact solution for color transparency^{*}

Jörg Hüfner^a and Boris Kopeliovich^{bc†}

^a Inst. für Theor. Physik der Universität , Philosophenweg 19, 69120 Heidelberg, Germany

^b Max-Planck Institut für Kernphysik, Postfach 103980, 69029 Heidelberg, Germany

^c Joint Institute for Nuclear Research, Dubna, 141980 Moscow Region, Russia

Abstract

A multichannel evolution equation is developed for the density matrix describing a hadronic wave packet produced by a virtual photon and propagation through a nuclear matter. This approach is dual to the quark-gluon representation, incorporates the effects of coherence and formation times and gives an exact solution for color transparency problem at any photon energy. We propose a procedure of data analysis, which provides an unambiguous way of detection of a color transparency signal even at medium energies.

1 Introduction

Exclusive electroproduction of vector mesons was suggested in [1] as an effective tool in search for color transparency (CT) [2]. The key idea is based upon absence of strict correlation between the photon's energy and virtuality, typical for reactions of quasielastic scattering. Data from the E665 experiment [3] nicely confirms the predicted value of the effect [1]. The statistical confidence of the observed growth of nuclear transparency with the photon virtuality Q^2 is, however, quite modest and new experiments are planned at lower energies (HERMES, TJNAF, ELFE).

Although CT phenomenon is most naturally interpreted in the quark-gluon representation, the calculations are easy only at energies $\nu \gg Q^2 R_A/2$, when size of the photon fluctuations is frozen during propagation through the nucleus. At lower energies the produced colorless wave packet is developing while it propagates through the nucleus. Accordingly, absorption in nuclear

^{*} A talk presented by B.K. at the second ELFE Workshop, September 22-27, 1996, Saint Malo, France

[†]Partially supported by the European Network: Hadronic Physics with Electromagnetic Probes. Contract No. FMRX-CT96-0008

matter varies and the expected CT effect may be substantially reduced. Such an evolution is controlled by the so-called formation time t_f ,

$$\frac{2\nu}{Q^2} < t_f < \frac{2\nu}{m_{V'}^2 - m_V^2} . \quad (1)$$

The bottom limit corresponds to most quickly expanding states of small size, $r_T^2 \propto 1/Q^2$. In order to observe a full effect of CT, one should make this time, $t_f^{min} \approx 1/m_N x_B \gg R_A$, where x_B is the Bjorken variable. The upper limit in (1) corresponds to a long evolution of a rather large-size wave packet consisted mostly from the two lightest states, V and V' . This time t_f^{max} controls the onset of CT.

There is another phenomenon, which follows from the quantum-mechanical uncertainty for the production time of the final wave packet, which is usually called coherence time,

$$t_c \approx \frac{2\nu}{Q^2 + m_V^2} . \quad (2)$$

This uncertainty may be interpreted as a lifetime of hadronic fluctuations of the photon. If this time substantially exceeds the nuclear size, $t_c \gg R_A$, one deals with a virtual hadronic, rather than with photonic beam. Correspondingly, nuclear attenuation increases [5, 6, 7].

Two methods for correct calculation of the wave packet evolution are known. One was developed in [4] using the quark-gluon representation and Feynman path integral technique. However, as far as the coherence time effects are involved, the path integrals approach becomes unreasonably complicate, and no solution is still found.

Another approach uses the hadronic basis for the photon fluctuations, which is dual to the quark-gluon basis, provided that completeness takes place. An exact solution, incorporating both coherence and formation time effects is found and the results are presented in this talk.

2 Coherence time

Coherence (interference) of the vector meson waves produced at different longitudinal coordinates is important both for coherent (the nucleus remains intact) and incoherent (the nucleus breaks up) electroproduction. Effect of coherence time exists even in Glauber approximation,

although a correct formula for incoherent electroproduction was derived only recently [5]. Vector mesons produced at different points separated by longitudinal distance Δz have a relative phase shift $q_c \Delta z$, where $q_c = (Q^2 + m_V^2)/2\nu$ is the longitudinal momentum transfer in $\gamma^* N \rightarrow VN$, Q^2 and ν are the virtuality and energy of the photon, respectively. Taking this into account one arrives at the following expression [5] for nuclear transparency defined as $Tr = \sigma_A/A\sigma_N$,

$$\begin{aligned} Tr_{inc} = & \frac{\sigma_{tot}^{VN}}{2A\sigma_{el}^{VN}} (\sigma_{in}^{VN} - \sigma_{el}^{VN}) \int d^2b \int_{-\infty}^{\infty} dz_2 \rho(b, z_2) \int_{-\infty}^{z_2} dz_1 \rho(b, z_1) \\ & \times e^{iq_c(z_2 - z_1)} \exp \left[-\frac{1}{2} \sigma_{tot}^{VN} \int_{z_1}^{z_2} dz \rho(b, z) \right] \exp \left[-\sigma_{in}^{VN} \int_{z_2}^{\infty} dz \rho(b, z) \right] \\ & + \frac{1}{A\sigma_{in}^{VN}} \int d^2b \left[1 - e^{-\sigma_{in}^{VN} T(b)} \right] - Tr_{coh} , \end{aligned} \quad (3)$$

$$Tr_{coh} = \frac{(\sigma_{tot}^{VN})^2}{4A\sigma_{el}^{VN}} \int d^2b \left| \int_{-\infty}^{\infty} dz \rho(b, z) e^{iq_c z} \exp \left[-\frac{1}{2} \sigma_{tot}^{VN} \int_z^{\infty} dz' \rho(b, z') \right] \right|^2 , \quad (4)$$

where Tr_{coh} corresponds to the coherent case [8, 5, 6]. Tr_{inc} , in contrast to the coherent case [7], decreases with energy from $Tr_{inc} = \sigma_{in}^{VA}/A\sigma_{in}^{VN}$ ($q_c \gg 1/R_A$) down to $Tr_{inc} = \sigma_{qel}^{VA}/A\sigma_{el}^{VN}$ ($q_c \ll 1/R_A$). Numerical examples are presented in [6, 7, 5].

Variation of t_c may be caused either by its ν - or Q^2 -dependence. In the latter case t_c decreases with Q^2 and the nuclear transparency grows, what is usually expected to be a signature of CT [1]. Our results for incoherent electroproduction of ρ -meson on lead are shown by dashed curves in Fig. 1 (more examples are in [6, 7, 5]). The predicted Q^2 -dependence is so steep that makes it quite problematic to observe a signal of CT on such a background.

3 Formation time

Inclusion of excited states of the vector meson into the multiple scattering series is known as Gribov's inelastic corrections [9]. CT corresponds to a special tuning of these corrections, when the diagonal and off diagonal amplitudes cancel in final state interaction at high Q^2 . The amplitudes we use satisfy such a condition, since we calculate the photoproduction amplitudes projecting to the Q^2 -dependent $q\bar{q}$ component of the photon wave function.

In the case of incoherent production one has to sum over all final states of the nucleus. Therefore,

the the wave packet should be described by density matrix $P_{ij} = \sum_{A^*} |\psi_i\rangle |\psi_j\rangle^+$. Wave function $|\psi_i\rangle$, has components γ^* , V , V' , etc. The evolution equation for the density matrix reads [10],

$$i \frac{d}{dz} \hat{P} = \hat{Q} \hat{P} - \hat{P} \hat{Q}^+ - \frac{i}{2} \sigma_{tot}^{VN} (\hat{T} \hat{P} + \hat{P} \hat{T}^+) + i \sigma_{el}^{VN} \hat{T} \hat{P} \hat{T}^+ , \quad (5)$$

$$\hat{Q} = \begin{pmatrix} 0 & 0 & 0 & \dots \\ 0 & q & 0 & \dots \\ 0 & 0 & q' & \dots \\ \vdots & \vdots & \vdots & \dots \end{pmatrix} ; \quad \hat{T} = \begin{pmatrix} 0 & 0 & 0 & \dots \\ \lambda & 1 & \epsilon & \dots \\ \lambda R & \epsilon & r & \dots \\ \vdots & \vdots & \vdots & \dots \end{pmatrix} , \quad (6)$$

where $\hat{Q}$ and $\hat{T}$ are the $(n+1) \times (n+1)$ matrices, and n is the number of states involved into consideration. $q, q' \dots$ are the transferred longitudinal momenta, $q(q') = (m_{V(V')}^2 + Q^2)/2\nu$. For other parameters we use notations from [11], $r = \sigma_{tot}^{V'N}/\sigma_{tot}^{VN}$, $\epsilon = f(VN \rightarrow V'N)/f(VN \rightarrow VN)$ and $R = f(\gamma N \rightarrow V'N)/f(\gamma N \rightarrow VN)$. The value of parameter $\lambda = f(\gamma N \rightarrow V'N)/f(VN \rightarrow VN)$ is inessential, since it cancels in nuclear transparency. The boundary condition for the density matrix is $P_{ij}(z \rightarrow -\infty) = \delta_{i0}\delta_{j0}$. Note, eq. (5) reproduces (3) if $\epsilon = 0$ and eq. (4) if $\sigma_{el}^{VN} = 0$.

We calculated the Q^2 -dependence of nuclear transparency for incoherent electroproduction of ρ -meson on lead in two-channel approximation, using the parameters in (6) as in [7] (more examples, including coherent production, radial excitations and other flavours are in [7, 10]). The two channels should well reproduce the onset of CT, while at higher energies, $1/x_B \ll m_N R_A$ the full CT may develop only after inclusion the the higher excitation states. The results for different energies are shown by solid curves in Fig. 1. Although the growth of nuclear transparency is steeper than what we expect in Glauber approximation, the difference is too small to be used as a signature of CT. Even the present state of art of Glauber-model calculations leaves enough freedom to fit in such a narrow corridor in nuclear transparency.

At this point we would like to soften our pessimism and suggest a method for unambiguous detection of onset of CT. The key idea is quite straitforward [10]: as soon as the variation of the coherence time may mock the CT effects, one should fix t_c . This can by done by means of a special selection of events with different Q^2 and ν . Starting from minimal energy $\nu_{min} = t_c Q^2/2$




Figure 1: Q^2 -dependence of nuclear transparency for ρ -meson electroproduction on lead at photon energies $\nu = 5, 10, 20$ and 30 GeV. Dashed curves correspond to Glauber approximation, solid curves are calculated with the evolution equation eq. (5).

Figure 2: The same as in Fig. 1, but with fixed $t_c = 2\nu_{\min}/m_\rho^2$.

with real photoproduction one should increase both Q^2 and ν , while $t_c = \text{const}$, in accordance with (2). Our predictions for Q^2 -dependence of the lead transparency at different values of minimal energy (or t_c) are depicted in Fig. 2 in comparison with Q^2 -independent expectations of Glauber approximation.

Concluding, we have developed a multichannel approach to incoherent exclusive electroproduction of vector mesons off nuclei, which incorporates the effects of coherent and formation times, as well as CT. Variation of the coherence time with the photon energy and Q^2 causes substantial changes of the nuclear transparency and may mock onset of CT. We suggest such a mapping of ν and Q^2 values, which keeps the coherence time constant. This helps to single out an unambiguous signal of CT at medium energies.

References

- [1] B.Z. Kopeliovich, J. Nemchik, N.N. Nikolaev and B.G. Zakharov, Phys. Lett B324 (1994) 469; Phys. Lett. B309 (1993) 179

- [2] P. Jain, B. Pire and J. P. Ralston, hep-ph/9511333, to appear in Physics Reports.
- [3] E665 Collab., M.R. Adams et al., Phys. Rev. Lett. 74 (1995) 1525
- [4] B.Z. Kopeliovich and B.G. Zakharov, Phys. Rev. D44 (1991) 3466
- [5] J. Hüfner, B.Z. Kopeliovich and J. Nemchik, Phys. Lett. B383 (1996) 362
- [6] J. Hüfner, B.Z. Kopeliovich and J. Nemchik, in the Proceedings of ELFE Workshop, ed. S.D. Bass and P.A.M. Guichon, p. 337, Editions Frontieres, 1995 hep-ph/9511215
- [7] B.Z. Kopeliovich and J. Nemchik, MPIH-V41-1995, nucl-th/9511018
- [8] T.H. Bauer et al., Rev. Mod. Phys. 50 (1978) 261
- [9] V.N. Gribov, Sov. Phys. JETP, 29 (1969) 483
- [10] J. Hüfner and B.Z. Kopeliovich, "Towards study of color transparency with medium-energy electron beams", nucl-th/9607034, in Proc. of the '95 - '96 workshop on Future Physics at HERA, NIKHEF 96-026, ed. by G. van der Steenhoven, Amsterdam, 1996, p. 33
- [11] J. Hüfner and B.Z. Kopeliovich, Phys. Rev. Lett. 76 (1996) 192